

J. Eric Hibbs, D.D.S., F.A.G.D.
www.ParkCitiesDental.com
5600 W. Lovers Ln @ Tollway #216
Dallas TX 75209-4318 214/351-2311



Park Cities Dental
E-mail: Dr@Hibbs.com

Member:
American Dental Association,
American Academy Of Cosmetic Dentistry,
Academy of General Dentistry,
Texas Dental Association,
Dallas County Dental Society,
American Orthodontic Society,
Federation Dentaire Internationale,
Association of Military Surgeons of the United States,
Commander, U. S. Navy Dental Corps Reserves,
Veterans of Foreign Wars

For a white, healthy smile:
◆ Mercury-free fillings
◆ Teeth whitening
◆ Porcelain veneers
◆ Custom-crafted crowns & bridges
◆ Non-surgical gum care
◆ Dentures & Partial
◆ Complex cases

For your comfort:
◆ Gentle cleanings
◆ Personal attention
◆ Thorough sterilization
◆ Flavored nitrous oxide
◆ TV/Music and chair massage to relax you

For your convenience:
◆ On-time appointments
◆ Children welcome
◆ Free on-site parking
◆ 24-hour emergency care 214.832.0992
◆ Insurance accepted and filed for you
◆ Amex, Visa, Discover and MasterCard

INSTRUCTIONS FOR PATIENTS FOLLOWING ORAL SURGERY

THINGS TO EXPECT:

- Bleeding:** Bleeding or “oozing” for the first 12 to 24 hours is normal.
Swelling: Swelling is normal following a surgical procedure in the mouth. It should reach its maximum in 18 hours and should diminish by the fifth post-operative day.
Discomfort: The most discomfort that you should experience will occur for a few hours after the sensation returns to your mouth and will diminish over the next few days.

THINGS TO DO:

- Bleeding:** Place gauze over extraction sites and maintain pressure by biting for at least one hour. Repeat as needed. Keep head elevated, and rest. Do not suck or spit excessively.
NOTE: Some “oozing” and discoloration of saliva is normal. If bleeding persists, replace the gauze with clean folded gauze placed over the extraction site, and maintain the pressure until the bleeding stops.
Rinsing: 24 hours after the surgery, rinse with warm salt water three to five times per day for seven days.
Swelling: Place ice or cold compresses on the region of surgery for ten minutes every half-hour for the first eight to twelve hours.
NOTE: Ice bags or cold compresses should be used only on the day of surgery.
Smoking: Avoid smoking during the healing period.
Discomfort: Take medications as directed for pain. If discomfort is severe, do not wait for the pain to begin before taking the medications since the medicine will require 45 minutes to one hour to take effect.
Diet: A nutritious liquid or soft diet may be necessary for the first two days. Instant breakfast drinks are beneficial. A fluid intake of at least two quarts a day is needed.
Physical Activity: For the first 24 to 48 hours, one should REST. Avoid any strenuous activity and do not drive while taking narcotic medications

THE DAYS AFTER SURGERY:

1. Brush teeth carefully.
2. Use warm salt water as a mouth rinse three to five times per day.
3. If antibiotics are prescribed, be sure to take all that have been prescribed, as directed.
4. Use warm, moist heat on face for swelling, if any. Continue until the swelling subsides. A warm, wet washcloth or heating pad will suffice.
5. If sutures were used, return to the office in five to seven days for removal, or as directed.
6. Dry socket is a delayed healing response, which may occur during the second to fourth post-operative day. It is associated with a throbbing pain on the side of the face, which may seem to be directed up toward the ear. In mild cases, simply increasing the pain medication can control the symptoms. If this is unsuccessful, please call your doctor.

CONTACT THE DOCTOR IF:

1. Bleeding is excessive and cannot be controlled.
2. Discomfort is poorly controlled.
3. Swelling is excessive, spreading, or continuing to enlarge after 48 hours.
4. Allergic reactions to medications occur which are causing a generalized rash or itchiness.

AFTER HOURS BEEPER: 214/832-0992